



# Quantum Theory and the Electronic Structure of Atoms

## Chapter 7

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

### Properties of Waves

**Wavelength** ( $\lambda$ ) is the distance between identical points on successive waves.

**Amplitude** is the vertical distance from the midline of a wave to the peak or trough.

**Frequency** ( $\nu$ ) is the number of waves that pass through a particular point in 1 second (Hz = 1 cycle/s).


The speed ( $u$ ) of the wave =  $\lambda \times \nu$

Maxwell (1873), proposed that **visible light consists of electromagnetic waves.**

**Electromagnetic radiation** is the emission and transmission of energy in the form of electromagnetic waves.

Speed of light ( $c$ ) in vacuum =  $3.00 \times 10^8$  m/s

**All electromagnetic radiation**  
 $\lambda \times \nu = c$



A photon has a frequency of  $6.0 \times 10^4$  Hz. Convert this frequency into wavelength (nm). Does this frequency fall in the visible region?

$$\lambda \times \nu = c$$

$$\lambda = c/\nu$$

$$\lambda = 3.00 \times 10^8 \text{ m/s} / 6.0 \times 10^4 \text{ Hz}$$

$$\lambda = 5.0 \times 10^3 \text{ m}$$

$$\lambda = 5.0 \times 10^{12} \text{ nm}$$

### Mystery #1, "Heated Solids Problem" Solved by Planck in 1900

When solids are heated, they emit electromagnetic radiation over a wide range of wavelengths.

Radiant energy emitted by an object at a certain temperature depends on its wavelength.

**Energy (light) is emitted or absorbed in discrete units (quantum).**

$E = h \times \nu$   
Planck's constant ( $h$ )  
 $h = 6.63 \times 10^{-34} \text{ J}\cdot\text{s}$

## Mystery #2, "Photoelectric Effect" Solved by Einstein in 1905


- Light has both:
1. wave nature
  2. particle nature

**Photon** is a "particle" of light

$$h\nu = KE + W$$

$$KE = h\nu - W$$

where  $W$  is the work function and depends how strongly electrons are held in the metal



When copper is bombarded with high-energy electrons, X rays are emitted. Calculate the energy (in joules) associated with the photons if the wavelength of the X rays is 0.154 nm.


$$E = h \times \nu$$

$$E = h \times c / \lambda$$


$$E = 6.63 \times 10^{-34} \text{ (J}\cdot\text{s)} \times 3.00 \times 10^8 \text{ (m/s)} / 0.154 \times 10^{-9} \text{ (m)}$$

$$E = 1.29 \times 10^{-15} \text{ J}$$

8



9




## Bohr's Model of the Atom (1913)

1.  $e^-$  can only have specific (quantized) energy values
2. light is emitted as  $e^-$  moves from one energy level to a lower energy level


$$E_n = -R_H \left( \frac{1}{n^2} \right)$$

$n$  (principal quantum number) = 1, 2, 3, ...


$R_H$  (Rydberg constant) =  $2.18 \times 10^{-18} \text{ J}$



11



12



13

TABLE 7.1 The Various Series in Atomic Hydrogen Emission Spectrum			
Series	$n_f$	$n_i$	Spectrum Region
Lyman	1	2, 3, 4, ...	Ultraviolet
Balmer	2	3, 4, 5, ...	Visible and ultraviolet
Paschen	3	4, 5, 6, ...	Infrared
Brackett	4	5, 6, 7, ...	Infrared

14

Calculate the wavelength (in nm) of a photon emitted by a hydrogen atom when its electron drops from the  $n = 5$  state to the  $n = 3$  state.

$$E_{\text{photon}} = \Delta E = R_H \left( \frac{1}{n_f^2} - \frac{1}{n_i^2} \right)$$

$$E_{\text{photon}} = 2.18 \times 10^{-18} \text{ J} \times (1/25 - 1/9)$$

$$E_{\text{photon}} = \Delta E = -1.55 \times 10^{-19} \text{ J}$$

$$E_{\text{photon}} = h \times c / \lambda$$

$$\lambda = h \times c / E_{\text{photon}}$$

$$\lambda = 6.63 \times 10^{-34} \text{ (J}\cdot\text{s)} \times 3.00 \times 10^8 \text{ (m/s)} / 1.55 \times 10^{-19} \text{ J}$$

$$\lambda = 1280 \text{ nm}$$

15

Why is  $e^-$  energy quantized?

De Broglie (1924) reasoned that  $e^-$  is both particle and wave.

$$2\pi r = n\lambda \quad \lambda = \frac{h}{mu}$$

$u$  = velocity of  $e^-$

$m$  = mass of  $e^-$



16


What is the de Broglie wavelength (in nm) associated with a 2.5 g Ping-Pong ball traveling at 15.6 m/s?

$$\lambda = h/mu \quad h \text{ in J}\cdot\text{s} \quad m \text{ in kg} \quad u \text{ in (m/s)}$$

$$\lambda = 6.63 \times 10^{-34} / (2.5 \times 10^{-3} \times 15.6)$$

$$\lambda = 1.7 \times 10^{-32} \text{ m} = 1.7 \times 10^{-23} \text{ nm}$$

17



18


### Chemistry in Action: Electron Microscopy

$$\lambda_e = 0.004 \text{ nm}$$

Electron micrograph of a normal red blood cell and a sickled red blood cell from the same person



STM image of iron atoms on copper surface



19


### Schrodinger Wave Equation

In 1926 Schrodinger wrote an equation that described both the particle and wave nature of the  $e^-$

Wave function ( $\psi$ ) describes:

1. energy of  $e^-$  with a given  $\psi$
2. probability of finding  $e^-$  in a volume of space

Schrodinger's equation can only be solved exactly for the hydrogen atom. Must approximate its solution for multi-electron systems.



20


### Schrodinger Wave Equation

$\psi$  is a function of four numbers called **quantum numbers** ( $n, l, m_l, m_s$ )


principal quantum number  $n$

$$n = 1, 2, 3, 4, \dots$$

distance of  $e^-$  from the nucleus



21



22

### Schrodinger Wave Equation

**quantum numbers:** ( $n, l, m_l, m_s$ )

angular momentum quantum number  $l$


for a given value of  $n, l = 0, 1, 2, 3, \dots n-1$

- |                           |                   |
|---------------------------|-------------------|
| $n = 1, l = 0$            | $l = 0$ s orbital |
| $n = 2, l = 0$ or $1$     | $l = 1$ p orbital |
| $n = 3, l = 0, 1,$ or $2$ | $l = 2$ d orbital |
|                           | $l = 3$ f orbital |


Shape of the "volume" of space that the  $e^-$  occupies

23

$l = 0$  (s orbitals)



$l = 1$  (p orbitals)



24

$l = 2$  (d orbitals)

$d_{xy}$     $d_{yz}$     $d_{zx}$     $d_{x^2-y^2}$     $d_{z^2}$

25

### Schrodinger Wave Equation

**quantum numbers:**  $(n, l, m_l, m_s)$

magnetic quantum number  $m_l$

for a given value of  $l$   
 $m_l = -l, \dots, 0, \dots, +l$

if  $l = 1$  (p orbital),  $m_l = -1, 0, \text{ or } 1$   
 if  $l = 2$  (d orbital),  $m_l = -2, -1, 0, 1, \text{ or } 2$

orientation of the orbital in space

26

$m_l = -1, 0, \text{ or } 1$       3 orientations in space

$2p_x$        $2p_y$        $2p_z$

27

$m_l = -2, -1, 0, 1, \text{ or } 2$       5 orientations in space

$3d_{xy}$     $3d_{yz}$     $3d_{zx}$   
 $3d_{x^2-y^2}$     $3d_{z^2}$

28

### Schrodinger Wave Equation

$(n, l, m_l, m_s)$

spin quantum number  $m_s$

$m_s = +\frac{1}{2} \text{ or } -\frac{1}{2}$

$m_s = +\frac{1}{2}$        $m_s = -\frac{1}{2}$

29

### Schrodinger Wave Equation

**quantum numbers:**  $(n, l, m_l, m_s)$

Existence (and energy) of electron in atom is described by its **unique** wave function  $\psi$ .

**Pauli exclusion principle** - no two electrons in an atom can have the same four quantum numbers.

Each seat is uniquely identified (E, R12, S8)  
 Each seat can hold only one individual at a time

30

**TABLE 7.2** Relation Between Quantum Numbers and Atomic Orbitals

$n$	$l$	$m_l$	Number of Orbitals	Atomic Orbital Designations
1	0	0	1	1s
2	0	0	1	2s
	1	-1, 0, 1	3	2p <sub>x</sub> , 2p <sub>y</sub> , 2p <sub>z</sub>
3	0	0	1	3s
	1	-1, 0, 1	3	3p <sub>x</sub> , 3p <sub>y</sub> , 3p <sub>z</sub>
	2	-2, -1, 0, 1, 2	5	3d <sub>xy</sub> , 3d <sub>yz</sub> , 3d <sub>xz</sub> , 3d <sub>z<sup>2</sup></sub> , 3d <sub>x<sup>2</sup>-y<sup>2</sup></sub>
⋮	⋮	⋮	⋮	⋮

31

### Schrodinger Wave Equation

**quantum numbers:**  $(n, l, m_l, m_s)$

Shell – electrons with the same value of  $n$

Subshell – electrons with the same values of  $n$  and  $l$

Orbital – electrons with the same values of  $n, l$ , and  $m_l$

How many electrons can an orbital hold?

If  $n, l$ , and  $m_l$  are fixed, then  $m_s = \frac{1}{2}$  or  $-\frac{1}{2}$

$\psi = (n, l, m_l, \frac{1}{2})$  or  $\psi = (n, l, m_l, -\frac{1}{2})$

An orbital can hold 2 electrons


32

How many 2p orbitals are there in an atom?

$n=2$   
↓  
2p  
↑  
 $l=1$

If  $l = 1$ , then  $m_l = -1, 0, \text{ or } +1$

3 orbitals






How many electrons can be placed in the 3d subshell?

$n=3$   
↓  
3d  
↑  
 $l=2$

If  $l = 2$ , then  $m_l = -2, -1, 0, +1, \text{ or } +2$

5 orbitals which can hold a total of 10 e<sup>-</sup>

33



The most stable arrangement of electrons in subshells is the one with the greatest number of parallel spins (**Hund's rule**).

Energy ↑

Ne 1s<sup>2</sup>2s<sup>2</sup>2p<sup>6</sup>

37

Order of orbitals (filling) in multi-electron atom

1s < 2s < 2p < 3s < 3p < 4s < 3d < 4p < 5s < 4d < 5p < 6s < 4f < 5d < 6p < 7s < 6d < 7p

**Electron configuration** is how the electrons are distributed among the various atomic orbitals in an atom.

number of electrons in the orbital or subshell

principal quantum number  $n$

angular momentum quantum number  $l$

**Orbital diagram**

H  $\uparrow$   
1s<sup>1</sup>

39

What is the electron configuration of Mg?

Mg 12 electrons

1s < 2s < 2p < 3s < 3p < 4s

1s<sup>2</sup>2s<sup>2</sup>2p<sup>6</sup>3s<sup>2</sup>    2 + 2 + 6 + 2 = 12 electrons

Abbreviated as [Ne]3s<sup>2</sup>    [Ne] 1s<sup>2</sup>2s<sup>2</sup>2p<sup>6</sup>

What are the possible quantum numbers for the last (outermost) electron in Cl?

Cl 17 electrons    1s < 2s < 2p < 3s < 3p < 4s

1s<sup>2</sup>2s<sup>2</sup>2p<sup>6</sup>3s<sup>2</sup>3p<sup>5</sup>    2 + 2 + 6 + 2 + 5 = 17 electrons

Last electron added to 3p orbital

$n = 3$      $l = 1$      $m_l = -1, 0, \text{ or } +1$      $m_s = \frac{1}{2} \text{ or } -\frac{1}{2}$  40

Outermost subshell being filled with electrons

41

TABLE 7.3 The Ground-State Electron Configurations of the Elements<sup>a</sup>

Atomic Number	Symbol	Electron Configuration	Atomic Number	Symbol	Electron Configuration	Atomic Number	Symbol	Electron Configuration
1	H	1s <sup>1</sup>	38	Sr	[Kr]5s <sup>2</sup>	75	Re	[Xe]6s <sup>2</sup> 4f <sup>14</sup> 5d <sup>5</sup>
2	He	1s <sup>2</sup>	39	Y	[Kr]5s <sup>2</sup> 4d <sup>1</sup>	76	Os	[Xe]6s <sup>2</sup> 4f <sup>14</sup> 5d <sup>6</sup>
3	Li	[He]2s <sup>1</sup>	40	Zr	[Kr]5s <sup>2</sup> 4d <sup>2</sup>	77	Ir	[Xe]6s <sup>2</sup> 4f <sup>14</sup> 5d <sup>7</sup>
4	Be	[He]2s <sup>2</sup>	41	Nb	[Kr]5s <sup>2</sup> 4d <sup>4</sup>	78	Pt	[Xe]6s <sup>2</sup> 4f <sup>14</sup> 5d <sup>8</sup>
5	B	[He]2s <sup>2</sup> 2p <sup>1</sup>	42	Mo	[Kr]5s <sup>2</sup> 4d <sup>5</sup>	79	Au	[Xe]6s <sup>2</sup> 4f <sup>14</sup> 5d <sup>9</sup>
6	C	[He]2s <sup>2</sup> 2p <sup>2</sup>	43	Tc	[Kr]5s <sup>2</sup> 4d <sup>5</sup>	80	Hg	[Xe]6s <sup>2</sup> 4f <sup>14</sup> 5d <sup>10</sup>
7	N	[He]2s <sup>2</sup> 2p <sup>3</sup>	44	Ru	[Kr]5s <sup>2</sup> 4d <sup>6</sup>	81	Tl	[Xe]6s <sup>2</sup> 4f <sup>14</sup> 5d <sup>10</sup> 6p <sup>1</sup>
8	O	[He]2s <sup>2</sup> 2p <sup>4</sup>	45	Rh	[Kr]5s <sup>2</sup> 4d <sup>7</sup>	82	Pb	[Xe]6s <sup>2</sup> 4f <sup>14</sup> 5d <sup>10</sup> 6p <sup>2</sup>
9	F	[He]2s <sup>2</sup> 2p <sup>5</sup>	46	Pd	[Kr]4d <sup>10</sup>	83	Bi	[Xe]6s <sup>2</sup> 4f <sup>14</sup> 5d <sup>10</sup> 6p <sup>3</sup>
10	Ne	[He]2s <sup>2</sup> 2p <sup>6</sup>	47	Ag	[Kr]5s <sup>2</sup> 4d <sup>10</sup>	84	Po	[Xe]6s <sup>2</sup> 4f <sup>14</sup> 5d <sup>10</sup> 6p <sup>4</sup>
11	Na	[Ne]3s <sup>1</sup>	48	Cd	[Kr]5s <sup>2</sup> 4d <sup>10</sup>	85	At	[Xe]6s <sup>2</sup> 4f <sup>14</sup> 5d <sup>10</sup> 6p <sup>5</sup>
12	Mg	[Ne]3s <sup>2</sup>	49	In	[Kr]5s <sup>2</sup> 4d <sup>10</sup> 5p <sup>2</sup>	86	Rn	[Xe]6s <sup>2</sup> 4f <sup>14</sup> 5d <sup>10</sup> 6p <sup>6</sup>
13	Al	[Ne]3s <sup>2</sup> 3p <sup>1</sup>	50	Sn	[Kr]5s <sup>2</sup> 4d <sup>10</sup> 5p <sup>2</sup>	87	Fr	[Rn]7s <sup>1</sup>
14	Si	[Ne]3s <sup>2</sup> 3p <sup>2</sup>	51	Sb	[Kr]5s <sup>2</sup> 4d <sup>10</sup> 5p <sup>3</sup>	88	Ra	[Rn]7s <sup>2</sup>
15	P	[Ne]3s <sup>2</sup> 3p <sup>3</sup>	52	Te	[Kr]5s <sup>2</sup> 4d <sup>10</sup> 5p <sup>4</sup>	89	Ac	[Rn]7s <sup>2</sup> 6d <sup>1</sup>
16	S	[Ne]3s <sup>2</sup> 3p <sup>4</sup>	53	I	[Kr]5s <sup>2</sup> 4d <sup>10</sup> 5p <sup>5</sup>	90	Th	[Rn]7s <sup>2</sup> 6d <sup>2</sup>
17	Cl	[Ne]3s <sup>2</sup> 3p <sup>5</sup>	54	Xe	[Kr]5s <sup>2</sup> 4d <sup>10</sup> 5p <sup>6</sup>	91	Pa	[Rn]7s <sup>2</sup> 6d <sup>1</sup> 7s <sup>1</sup>
18	Ar	[Ne]3s <sup>2</sup> 3p <sup>6</sup>	55	Cs	[Xe]6s <sup>1</sup>	92	U	[Rn]7s <sup>2</sup> 7p <sup>1</sup> 6d <sup>1</sup>
19	K	[Ar]4s <sup>1</sup>	56	Ba	[Xe]6s <sup>2</sup>	93	Np	[Rn]7s <sup>2</sup> 6d <sup>1</sup> 7p <sup>1</sup>

42

