





Chemical Bonding I: Basic Concepts


Chapter 9


Li  *Handwritten notes: 1s² 2s¹*


Be  *Handwritten notes: 1s² 2s²*


B  *Handwritten notes: 1s² 2s² 2p¹*

C  *Handwritten notes: 1s² 2s² 2p²*

N  *Handwritten notes: 1s² 2s² 2p³*

O  *Handwritten notes: 1s² 2s² 2p⁴*

F  *Handwritten notes: 1s² 2s² 2p⁵*

Ne  *Handwritten notes: 1s² 2s² 2p⁶*

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

Valence electrons are the outer shell electrons of an atom. The valence electrons are the electrons that participate in chemical bonding.

Group	e ⁻ configuration	# of valence e ⁻
1A	ns ¹	1
2A	ns ²	2
3A	ns ² np ¹	3
4A	ns ² np ²	4
5A	ns ² np ³	5
6A	ns ² np ⁴	6
7A	ns ² np ⁵	7

2

Lewis Dot Symbols for the Representative Elements & Noble Gases

1A	2A	3A	4A	5A	6A	7A	8A
•H	•He•	•B•	•C••	•N•••	•O••••	•F•••••	•Ne••••••
•Li	•Be••	•Al•	•Si••	•P•••	•S••••	•Cl•••••	•Ar••••••
•Na	•Mg••	•Ga•	•Ge••	•As•••	•Se••••	•Br•••••	•Kr••••••
•K	•Ca••	•In•	•Sn••	•Sb•••	•Te••••	•I•••••	•Xe••••••
•Rb	•Sr••	•Tl•	•Pb••	•Bi•••	•Po••••	•At•••••	•Rn••••••
•Cs	•Ba••	•Tl•	•Pb••	•Bi•••	•Po••••	•At•••••	•Rn••••~•
•Fr	•Ra••						


3

The Ionic Bond

Ionic bond: the electrostatic force that holds ions together in an ionic compound.

$\bullet\text{Li} + \bullet\text{F}\bullet \longrightarrow \text{Li}^+ \text{F}^-$
 $1s^2 2s^1 1s^2 2s^2 2p^5 \rightarrow [1s^2] 2s^2 2p^6$

$\bullet\text{Li} \longrightarrow \text{Li}^+ + e^-$
 $e^- + \bullet\text{F}\bullet \longrightarrow \text{F}^-$
 $\text{Li}^+ + \text{F}^- \longrightarrow \text{Li}^+ \text{F}^-$



4

Electrostatic (Lattice) Energy

Lattice energy (U) is the energy required to completely separate one mole of a solid ionic compound into gaseous ions.


E is the potential energy
 Q_+ is the charge on the cation
 Q_- is the charge on the anion
 r is the distance between the ions

$$E = k \frac{Q_+ Q_-}{r}$$

Lattice energy increases as **Q increases** and/or as **r decreases**.

Compound	Lattice Energy (kJ/mol)	Q: +2,-1	Q: +2,-2
MgF ₂	2957	Q: +2,-1	
MgO	3938	Q: +2,-2	
LiF	1036		$r\text{F}^- < r\text{Cl}^-$
LiCl	853		

5




Compound	Lattice Energy (kJ/mol)	Melting Point (°C)
LiF	1017	845
LiCl	828	610
LiBr	787	550
LiI	732	450
NaCl	788	801
NaBr	736	750
NaI	686	662
KCl	699	772
KBr	689	735
KI	632	680
MgCl ₂	2527	714
Na ₂ O	2570	Sub ^a
MgO	3890	2800

^aNa₂O sublimes at 1275°C.

7

Chemistry In Action: Sodium Chloride: A Common and Important Compound



Mining Salt




Solar Evaporation for Salt




8

A **covalent bond** is a chemical bond in which two or more electrons are shared by two atoms.

Why should two atoms share electrons?




Lewis structure of F₂




9


Lewis structure of water



Double bond – two atoms share two pairs of electrons



Triple bond – two atoms share three pairs of electrons



10

Lengths of Covalent Bonds




TABLE 9.2
Average Bond Lengths of
Some Common Single,
Double, and Triple Bonds

Bond Type	Bond Length (pm)
C-H	107
C-O	143
C-O	121
C-C	154
C-C	133
C=C	130
C-N	112
C=N	138
C=N	116
N-O	136
N-O	122
O-H	96

Bond Lengths

Triple bond < Double Bond < Single Bond

11

TABLE 9.3 Comparison of Some General Properties of an Ionic Compound and a Covalent Compound

Property	NaCl	CCl ₄
Appearance	White solid	Colorless liquid
Melting point (°C)	801	-23
Molar heat of fusion* (kJ/mol)	30.2	2.5
Boiling point (°C)	1413	76.5
Molar heat of vaporization* (kJ/mol)	600	30
Density (g/cm ³)	2.17	1.59
Solubility in water	High	Very low
Electrical conductivity		
Solid	Poor	Poor
Liquid	Good	Poor

*Molar heat of fusion and molar heat of vaporization are the amounts of heat needed to melt 1 mole of the solid and to vaporize 1 mole of the liquid, respectively.

12

Polar covalent bond or **polar bond** is a covalent bond with greater electron density around one of the two atoms

electron poor region electron rich region

e⁻ poor e⁻ rich

H — F

δ^+ δ^-

13

Electronegativity is the ability of an atom to attract toward itself the electrons in a chemical bond.

Electron Affinity - **measurable**, Cl is highest

$$X_{(g)} + e^- \longrightarrow X^-_{(g)}$$



Electronegativity - **relative**, F is highest

13

The Electronegativities of Common Elements

Increasing electronegativity																							
Increasing electronegativity																							
1A																	5A	4A	3A	2A	6A	7A	8A
H 2.1																	B 2.0	C 2.5	N 3.0	O 3.5	F 4.0	Ne	
Li 1.0	Be 1.5															Al 1.5	Si 1.8	P 2.1	S 2.5	Cl 3.0	Ar		
Na 0.9	Mg 1.2	3B	4B	5B	6B	7B	8B	9B	10B	11B	12B	Ga 1.7	Ge 1.8	As 2.0	Se 2.4	Br 2.8	Kr						
K 0.8	Ca 1.0	Sc	Ti 1.3	V 1.6	Cr 1.6	Mn 1.5	Fe 1.8	Cu 1.9	Zn 1.6	2B	3B	4B	5B	6B	7B	8B	9B						
Rb 0.8	Sr 1.0	Y	Zr 1.4	Nb 1.6	Mo 2.0	Tc 1.9	Ru 2.2	Rh 2.2	Pd 2.2	Ag 1.9	Cd 1.7	In 1.7	Sn 1.8	Sb 1.9	Te 2.1	I 2.5	Xe						
Cs 0.7	Ba 0.9	La-Lu	Hf 1.3	Ta 1.5	W 1.7	Re 1.9	Os 2.2	Ir 2.2	Pt 2.2	Au 2.4	Hg 1.9	Tl 1.8	Pb 1.8	Bi 1.9	Po 2.0	At 2.2	Rn						
Fr 0.7	Ra 0.9																	17					

15



Classify the following bonds as ionic, polar covalent, or covalent: The bond in CsCl; the bond in H₂S; and the NN bond in H₂NNH₂.

Cs - 0.7	Cl - 3.0	3.0 - 0.7 = 2.3	Ionic
H - 2.1	S - 2.5	2.5 - 2.1 = 0.4	Polar Covalent
N - 3.0	N - 3.0	3.0 - 3.0 = 0	Covalent

18



Writing Lewis Structures

1. Draw skeletal structure of compound showing what atoms are bonded to each other. Put least electronegative element in the center.
2. Count total number of valence e⁻. Add 1 for each negative charge. Subtract 1 for each positive charge.
3. Complete an octet for all atoms **except** hydrogen
4. If structure contains too many electrons, form double and triple bonds on central atom as needed.

19

Write the Lewis structure of nitrogen trifluoride (NF₃).


- Step 1 – N is less electronegative than F, put N in center
- Step 2 – Count valence electrons N - 5 (2s²2p³) and F - 7 (2s²2p⁵)
- $$5 + (3 \times 7) = 26 \text{ valence electrons}$$
- Step 3 – Draw single bonds between N and F atoms and complete octets on N and F atoms.
- Step 4 - Check, are # of e⁻ in structure equal to number of valence e⁻?
- $$3 \text{ single bonds } (3 \times 2) + 10 \text{ lone pairs } (10 \times 2) = 26 \text{ valence electrons}$$



20

Write the Lewis structure of the carbonate ion (CO₃²⁻).

- Step 1 – C is less electronegative than O, put C in center
- Step 2 – Count valence electrons C - 4 (2s²2p²) and O - 6 (2s²2p⁴)
- $$-2 \text{ charge} - 2e^-$$
- $$4 + (3 \times 6) + 2 = 24 \text{ valence electrons}$$
- Step 3 – Draw single bonds between C and O atoms and complete octet on C and O atoms.
- Step 4 - Check, are # of e⁻ in structure equal to number of valence e⁻?
- $$3 \text{ single bonds } (3 \times 2) + 10 \text{ lone pairs } (10 \times 2) = 26 \text{ valence electrons}$$
- Step 5 - Too many electrons, form double bond and re-check # of e⁻





$$\begin{aligned} 2 \text{ single bonds } (2 \times 2) &= 4 \\ 1 \text{ double bond} &= 4 \\ 8 \text{ lone pairs } (8 \times 2) &= 16 \\ \hline \text{Total} &= 24 \end{aligned}$$



21

Two possible skeletal structures of formaldehyde (CH₂O)




An atom's **formal charge** is the difference between the number of valence electrons in an isolated atom and the number of electrons assigned to that atom in a Lewis structure.

$$\text{formal charge on an atom in a Lewis structure} = \text{total number of valence electrons in the free atom} - \text{total number of nonbonding electrons} - \frac{1}{2} \left(\text{total number of bonding electrons} \right)$$

The sum of the formal charges of the atoms in a molecule or ion must equal the charge on the molecule or ion.

22




$$\begin{aligned} \text{C} - 4 e^- & & 2 \text{ single bonds } (2 \times 2) &= 4 \\ \text{O} - 6 e^- & & 1 \text{ double bond} &= 4 \\ \hline 2\text{H} - 2 \times 1 e^- & & 2 \text{ lone pairs } (2 \times 2) &= 4 \\ 12 e^- & & \text{Total} &= 12 \end{aligned}$$

$$\text{formal charge on an atom in a Lewis structure} = \text{total number of valence electrons in the free atom} - \text{total number of nonbonding electrons} - \frac{1}{2} \left(\text{total number of bonding electrons} \right)$$

$$\text{formal charge on C} = 4 - 2 - \frac{1}{2} \times 6 = -1$$

$$\text{formal charge on O} = 6 - 2 - \frac{1}{2} \times 6 = +1$$

23



$$\begin{aligned} \text{C} - 4 e^- & & 2 \text{ single bonds } (2 \times 2) &= 4 \\ \text{O} - 6 e^- & & 1 \text{ double bond} &= 4 \\ \hline 2\text{H} - 2 \times 1 e^- & & 2 \text{ lone pairs } (2 \times 2) &= 4 \\ 12 e^- & & \text{Total} &= 12 \end{aligned}$$

$$\text{formal charge on an atom in a Lewis structure} = \text{total number of valence electrons in the free atom} - \text{total number of nonbonding electrons} - \frac{1}{2} \left(\text{total number of bonding electrons} \right)$$

$$\text{formal charge on C} = 4 - 0 - \frac{1}{2} \times 8 = 0$$

$$\text{formal charge on O} = 6 - 4 - \frac{1}{2} \times 4 = 0$$





24

Formal Charge and Lewis Structures


- For neutral molecules, a Lewis structure in which there are no formal charges is preferable to one in which formal charges are present.
- Lewis structures with large formal charges are less plausible than those with small formal charges.
- Among Lewis structures having similar distributions of formal charges, the most plausible structure is the one in which negative formal charges are placed on the more electronegative atoms.

Which is the most likely Lewis structure for CH_2O ?





25

A **resonance structure** is one of two or more Lewis structures for a single molecule that cannot be represented accurately by only one Lewis structure.





What are the resonance structures of the carbonate (CO_3^{2-}) ion?



26

Exceptions to the Octet Rule


The Incomplete Octet




27

Exceptions to the Octet Rule

Odd-Electron Molecules





The Expanded Octet (central atom with principal quantum number $n > 2$)








28

Chemistry In Action: Just Say NO



The enthalpy change required to break a particular bond in one mole of gaseous molecules is the **bond enthalpy**.

Bond Enthalpy





Bond Enthalpies

Single bond < Double bond < Triple bond

30

Average **bond enthalpy** in polyatomic molecules



$$\text{Average OH bond enthalpy} = \frac{502 + 427}{2} = 464 \text{ kJ}$$

TABLE 4 Bond Bond Enthalpies of Diatomic Molecules and Average Bond Enthalpies for Bonds in Polyatomic Molecules


Bond	Bond Enthalpy (kJ/mol)	Bond	Bond Enthalpy (kJ/mol)
H-H	436	C-S	259
H-N	391	C=O	743
H-O	463	C≡N	891
H-F	568	C≡S	414
H-Cl	431	C≡N	891
H-Br	366	C≡O	1072
H-I	297	C≡O	1072
F-F	158	C≡O	1072
F-Cl	427	C≡O	1072
F-Br	253	C≡O	1072
F-I	175	C≡O	1072
Cl-Cl	242	C≡O	1072
Cl-Br	205	C≡O	1072
Cl-I	168	C≡O	1072
Br-Br	193	C≡O	1072
Br-I	147	C≡O	1072
I-I	151	C≡O	1072
C-C	347	C≡O	1072
C=C	614	C≡O	1072
C≡C	839	C≡O	1072
C-O	358	C≡O	1072
C=O	743	C≡O	1072
C-N	305	C≡O	1072
C=N	630	C≡O	1072
C≡N	891	C≡O	1072
C-S	259	C≡O	1072
C-Se	226	C≡O	1072
C-Te	208	C≡O	1072
C-P	306	C≡O	1072
C-As	237	C≡O	1072
C-Sb	226	C≡O	1072
C-Bi	203	C≡O	1072
C-H	414	C≡O	1072
C-D	411	C≡O	1072
C-T	408	C≡O	1072
C-F	485	C≡O	1072
C-Cl	339	C≡O	1072
C-Br	276	C≡O	1072
C-I	213	C≡O	1072
C-O	358	C≡O	1072
C-S	259	C≡O	1072
C-N	305	C≡O	1072
C-P	306	C≡O	1072
C-B	389	C≡O	1072
C-Al	325	C≡O	1072
C-Ga	293	C≡O	1072
C-In	265	C≡O	1072
C-Tl	233	C≡O	1072
C-Si	318	C≡O	1072
C-Ge	286	C≡O	1072
C-Sn	255	C≡O	1072
C-Pb	223	C≡O	1072
C-Bi	203	C≡O	1072
C-Po	182	C≡O	1072

31



Bond Enthalpies (BE) and Enthalpy changes in reactions

Imagine reaction proceeding by breaking all bonds in the reactants and then using the gaseous atoms to form all the bonds in the products.

$$\Delta H^\circ = \text{total energy input} - \text{total energy released} \\ = \Sigma \text{BE}(\text{reactants}) - \Sigma \text{BE}(\text{products})$$




32



33

Use bond enthalpies to calculate the enthalpy change for:



$$\Delta H^\circ = \Sigma \text{BE}(\text{reactants}) - \Sigma \text{BE}(\text{products})$$

Type of bonds broken	Number of bonds broken	Bond enthalpy (kJ/mol)	Enthalpy change (kJ/mol)
H—H	1	436.4	436.4
F—F	1	156.9	156.9
Type of bonds formed	Number of bonds formed	Bond enthalpy (kJ/mol)	Enthalpy change (kJ/mol)
H—F	2	568.2	1136.4

$$\Delta H^\circ = 436.4 + 156.9 - 2 \times 568.2 = -543.1 \text{ kJ/mol}$$

34